

**\*\*REPRESENTATIVE DATASHEET\*\*****Goat anti-human Factor XI (FXI)**

Peroxidase conjugated IgG

0.2 mg

**Product #:** GAFXI-HRP**Lot #:** XXXX**Expiry date:** XXXX

Store at -10 to -20°C

1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5  
905-304-9896 • 800-903-6020 • fax 905-304-9897

For Research Use Only.

Not for use in diagnostic procedures.

**Description of Factor XI (FXI)**

Factor XI (FXI, plasma thromboplastin antecedent) is a coagulation protein produced in the liver that circulates in plasma at approximately 5 µg/ml (30 nM). The mass of FXI is 160 kDa as determined by SDS-PAGE under non-reducing conditions and 80 kDa upon reduction. FXI consists of two identical 80 kDa subunits linked by disulphide bonds. Each subunit consists of a tandem repeat of four apple domains followed by a serine protease catalytic domain. Cleavage of FXI by activated factor XII or thrombin converts each subunit into a two-chain form and generates two active sites per FXIa molecule. The mass of FXIa is 160 kDa unreduced, but upon reduction FXIa migrates as a heavy chain of 50 kDa and a light chain of 30 kDa. The catalytic site of FXIa resides in the light chain. In plasma, FXI or FXIa circulates in non-covalent 1:1 complex with high molecular weight kininogen, which acts as a cofactor in the activation of FXI by activated factor XII. The activity of FXIa is regulated by platelets and by several proteinase inhibitors including, in order of decreasing importance, C1-inhibitor,  $\alpha_2$ antiplasmin,  $\alpha_1$ antitrypsin and antithrombin. Heparin has relatively little effect on the rate of inhibition of FXIa by antithrombin. The only known natural substrate for activated FXI (FXIa) is factor IX (Christmas factor) and the only cofactor required for this reaction is ionized calcium<sup>1-3</sup>.

**REFERENCES and REVIEWS**

1. Willemin WA, Minnema M, Meijers JCM, Roem D, Erenberg AJM, Nuijens JH, ten Cate H, Hack EC; Inactivation of Factor XIa in Human Plasma Assessed by Measuring Factor XIa-Protease Inhibitor Complexes: Major Role for C1-Inhibitor. *Blood* 85:1517, 1995.
2. DeLa Cadena R, Watchfogel YT, Colman RW, in Hemostasis and Thrombosis, 3<sup>rd</sup> Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 219-240, J.B. Lippincott Co., Philadelphia, 1994.
3. Baglia FA, Seaman FS, Walsh, PN; The Apple 1 and 4 domains of Factor XI Act to Synergistically Promote the Surface-Mediated Activation of Factor XI by Factor XIIa. *Blood* 85:2078, 1995.

**Product Specifications****Description:**

Vial containing XXXX ml of whole IgG conjugated to horseradish peroxidase (HRP) through carbohydrate groups. Total protein is 0.2 mg.

**Format:**

IgG-HRP conjugate as a clear, slightly red-brown liquid.

**Host Animal:**

Goat

**Immunogen:**

Human Factor XI purified from plasma.

**Concentration:**IgG-HRP concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ( $E_{1\%}^{1\text{cm}}$ ) of 14.**Buffer:**

A buffered stabilizer solution containing 50% (v/v) glycerol.

**Storage:**

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use. Avoid exposure to sodium azide as this is an inhibitor of peroxidase activity.

**Specificity:**

Prior to conjugation, this antibody was specific for FXI as demonstrated by immunoelectrophoresis and ELISA.

**Applications:**

Suitable as a source of peroxidase-labeled antibodies to FXI.

**Rz Ratio (Reinheitszahl,  $A_{403}/A_{280}$ ):**

XXXX as determined spectrophotometrically.

**Related Products:**

Cat #: GAFXI-IG	Goat anti-human FXI, IgG from antiserum
Cat #: GAFXI-AP	Goat anti-human FXI, affinity purified IgG
Cat #: SAFXI-IG	Sheep anti-human FXI, IgG from antiserum
Cat #: SAFXI-AP	Sheep anti-human FXI, affinity purified IgG
Cat #: FXI-EIA	Paired antibody set for ELISA of FXI, 5 x 96 wells
Cat #: FXI-DP	Human plasma deficient in FXI, immune depleted
Cat #: FXI-AG	VisuLize FXI Antigen ELISA Kit

Visit our site ([www.affinitybiologicals.com](http://www.affinitybiologicals.com)) for details.

**Limited Warranty:** This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:  
**AFFINITY BIOLOGICALS INC.**  
1348 Sandhill Drive  
Ancaster ON CANADA L9G 4V5  
Tel: (905) 304-9896  
(800) 903-6020  
Fax: (905) 304-9897  
[info@affinitybiologicals.com](mailto:info@affinitybiologicals.com)