

## \*\*REPRESENTATIVE DATASHEET\*\*

# Sheep anti-human Antithrombin (ATIII)

Peroxidase Conjugated Affinity-Purified IgG 0.1 mg

Product #:SAAT-APHRPLot #:XXXXExpiry date:XXXX

Store at -10 to -20°C.

#### 1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5 905-304-9896 \* 800-903-6020 \* fax 905-304-9897

For Research Use Only. Not for use in diagnostic procedures.

## Description of Antithrombin (ATIII)

Antithrombin, also known as Antithrombin III (ATIII), is a member of the SERPIN family of proteinase inhibitors and the primary inhibitor of thrombin in plasma. It is produced in the liver and circulates in plasma at ~200 µg/ml (~3.5 µM). Antithrombin inhibits a broad spectrum of serine proteases including thrombin, activated forms of factor X, factor IX, factor XI, factor XII, as well as kallikrein, plasmin and urokinase. Enzyme inhibition by antithrombin occurs through proteolytic cleavage at Arg<sup>385</sup>-Ser<sup>386</sup> and subsequent rapid formation of a stable, inactive 1:1 enzyme-antithrombin complex. Heparin has a profound accelerating effect on the inhibitory activity of antithrombin towards some enzymes. For example, the rate of inhibition of thrombin and activated factor X is increased 1000-fold in the presence of optimal concentrations of heparin, whereas heparin has relatively little effect on the inhibition rate of activated factor XI, activated factor XII and kallikrein. Antithrombin is a single chain molecule with a molecular weight of 59 kDa. Interaction with thrombin results in an SDS-stable thrombin-antithrombin complex of 96 kDa1-3.

## **REFERENCES** and **REVIEWS**

**1**. Damus PS, Rosenberg RD; Antithrombin – Heparin Cofactor; Methods in Enzymology <u>45</u>, pp 653-669, 1976.

**2.** Harpel PC; Blood Proteolytic Enzyme Inhibitors: Their Role in Modulating Blood Coagulation and Fibrinolytic Enzyme Pathways; in Hemostasis and Thrombosis, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 738-747, J.B. Lippincott Co., Philadelphia PA, USA, 1982.

**3.** Griffith MJ, Noyes CM, Church FC; Reactive Site Peptide Structural Similarity between Heparin Cofactor II and Antithrombin III; JBC:260, pp 2218-2225, 1985.

#### **Description:**

Vial containing XXXX ml of affinity-purified IgG conjugated to horseradish peroxidase (HRP) through carbohydrate groups. Total protein is 0.1 mg.

#### Format:

APIgG-HRP conjugate as a clear, slightly red-brown liquid.

#### Host Animal:

Sheep

#### Immunogen:

Human antithrombin purified from plasma.

### Concentration:

APIgG-HRP concentration is **XXXX** mg/ml, determined by absorbance using an extinction coefficient ( $E^{1\%}_{280}$ ) of 14.

#### Buffer:

A buffered stabilizer solution containing 50% (v/v) glycerol.

#### Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use. Avoid exposure to sodium azide as this is an inhibitor of peroxidase activity.

#### Specificity:

Prior to conjugation this antibody was specific for antithrombin as demonstrated by immunoelectrophoresis and ELISA.

#### Applications:

Suitable as a source of peroxidase-labeled antibodies to ATIII.

### Rz Ratio (Reinheitszahl, A403/A280):

XXXX as determined spectrophotometrically.

#### Related Products:

Cat #: GAAT-IG	Goat anti- ATIII, whole IgG from antiserum
Cat #: GAAT-AP	Goat anti-ATIII, affinity-purified IgG
Cat #: SAAT-IG	Sheep anti-ATIII, whole IgG from antiserum
Cat #: SAAT-AP	Sheep anti-ATIII, affinity-purified lgG
Cat #: ATIII-EIA	Paired antibody set for ELISA of ATIII, 5 x 96 wells
Cat #: TAT-EIA	Paired antibody ELISA of Thrombin-ATIII complexes
Cat #: AT-NU10	Paired antibody set for ATIII neutralization in plasma
Cat #: ATIII-DP	Human plasma deficient in ATIII, immune depleted
Cat #: ATHC-DP	Human ATIII & HCII def plasma, immune depleted

Visit our site (www.affinitybiologicals.com) for details.

**Limited Warranty:** This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantibility or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by: **AFFINITY BIOLOGICALS INC.** 1348 Sandhill Drive Ancaster ON CANADA L9G 4V5 Tel: (905) 304-9896 (800) 903-6020 Fax: (905) 304-9897 info@affinitybiologicals.com