

REPRESENTATIVE DATASHEET

Sheep anti-human Factor V (FV)

Whole IgG from antiserum 10 mg

Product #: SAFV-IG

Lot #: XXXX Expiry date: XXXX

Store at -10 to -20°C

1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5 905·304·9896 • 800·903·6020 • fax 905·304·9897

For Research Use Only. Not for use in diagnostic procedures.

Description of Factor V (FV)

Factor V (formerly referred to as accelerator globulin and labile factor) is a large glycoprotein (320 kDa) that is produced in the liver. The gene that encodes factor V (FV) is located on chromosome 1. A congenital deficiency of FV is a hemorrhagic disorder inherited as an autosomal recessive disease.

The concentration of FV in plasma is typically 10 μ g/ml. FV is a pro-cofactor that is activated through limited proteolysis by thrombin, or by activated factor X in the presence of phospholipid surface. Other physiologic activators of FV include plasmin, neutrophil elastase and platelet calpain. The activated cofactor (FVa) is an essential component of the prothrombin activator complex, which consists of FVa, activated factor X, calcium and anionic phospholipid surface. The intact prothrombinase complex activates prothrombin to thrombin at a rate 300,000-fold greater than activated factor X alone. In a positive feedback loop, the thrombin generated accelerates its own generation by activating more FV to FVa. Thrombin also acts to down-regulate FVa indirectly by activating Protein C, which inactivates FVa cofactor activity¹⁻³.

REFERENCES and REVIEWS

- 1. Kane WH, Davie EW; Blood Coagulation Factors V and VIII: Structural and functional similarities and their relationship to hemorrhagic and thrombotic disorders. Blood 71:539, 1988.
- 2. Hoyer, LW, Wyshock EG, Colman RW, in Hemostasis and Thrombosis, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 109-133, J.B. Lippincott Co., Philadelphia, 1994.
- 3. Nesheim ME, Katzmann JA, Tracy PB, Mann KG; in Methods in Enzymology 80:249, 1980.

Product Specifications

Description:

Vial containing XXXX ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 10 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Sheep

Immunogen:

Human factor V purified from plasma.

Concentration:

IgG concentration is $\frac{XXXX}{M}$ mg/ml, determined by absorbance using an extinction coefficient ($E^{1\%}_{280}$) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20° C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for factor V as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to human factor V.

Neutralizing activity:

One Bethesda Units/ml IgG against normal plasma.

One Bethesda unit/ml is defined as the amount of inhibitor that resulted in 50% residual FV activity after 2 hours at 37°C (Kasper CK et al, Thromb Diath Haemorrh **34**:869, 1975).

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Ī	I I	WW	N/	WW	D-4-	VVVV
	Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
	Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Cat #: SAFV-AP
Cat #: SAFV-HRP
Cat #: FV-EIA
Cat #: FV-DP
Sheep anti-human FV, affinity purified IgG
Sheep anti-human FV, IgG-peroxidase conjugate
Paired antibody set for ELISA of FV, 5 x 96 wells
Human plasma deficient in FV, immune depleted

Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantibility or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by: **AFFINITY BIOLOGICALS INC.**

AFFINITY BIOLOGICALS INC. 1348 Sandhill Drive

Ancaster ON CANADA L9G 4V5

Tel: (905) 304-9896 (800) 903-6020 Fax: (905) 304-9897

info@affinitybiologicals.com