

REPRESENTATIVE DATASHEET

Sheep anti-rabbit Thrombin

Affinity-Purified IgG 0.5 mg

Product #:	SART-AP	
Lot #:	XXXX	
Expiry date:	XXXX	

Store at -10 to -20°C

1348 Sandhill Drive. Ancaster, Ontario, Canada L9G 4V5 905-304-9896 * 800-903-6020 * fax 905-304-9897

For Research Use Only. Not for use in diagnostic procedures.

Description of Thrombin

Thrombin (EC3.4.21.5, α-thrombin) is the product of proteolytic activation of the zymogen prothrombin. Human thrombin is a two-chain serine protease with a mass of 37 kDa. The active site is located within the heavy chain. Thrombin has a high specificity for certain arginine bonds in protein substrates. The primary substrate is fibrinogen which thrombin converts to fibrin through the cleavage of four arginyl-glycyl peptide bonds. Thrombin is also an important activator of platelets, factor XIII, Protein C and TAFI (Plasma procarboxypeptidase B). In a positive feedback mechanism, thrombin increases the rate of its own production by activation of factors VIII and V. The rate of thrombin production is subsequently limited indirectly through the activation of Protein C by thrombin, which then inactivates the activated cofactors VIII and V. The binding of thrombin to thrombomodulin on the cell surface dramatically alters thrombin's specificity, increasing it's activity toward Protein C and TAFI, and decreasing it's activity toward fibrinogen and activating cofactors VIII and V. In plasma, thrombin activity is inhibited primarily by antithrombin and to a lesser extent heparin cofactor II. The rate of inhibition by both of these inhibitors is profoundly increased in the presence of optimal concentrations of heparin. Other physiological inhibitors of thrombin in the absence of heparin include α_2 -macroglobulin and α_1 -antitrypsin¹⁻⁴.

REFERENCES and **REVIEWS**

Document OPI0181, rev 2

1. Mann KG; Prothrombin and Thrombin; in Hemostasis and Thrombosis, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 184-199, J.B. Lippincott Co., Philadelphia PA, USA, 1994.

2. Stubbs MT, Bode W; A Player of Many Parts: The Spotlight Falls on Thrombin's Structure; Thrombosis Research 69, pp 1-58, 1993.

3. Downing MW, Bloom JW, Mann KG; Comparison of the Inhibition of Thrombin by Three Plasma Protease Inhibitors; Biochemistry 17, pp 2649-2653, 1978.

4. Weitz JI, Hudoba M, Massel D, Maranganore J, Hirsh J; Clot-bound Thrombin is Protected from Inhibition by Heparin-ATIII but is Susceptible to -ATIII independent Inhibitors; J. Clin. Invest 86, pp 385-391, 1990.

Product Specifications

Description:

Vial containing XXXX ml of IgG purified by chromatography on immobilized PPACK-thrombin. Total protein is 0.5 mg.

Format:

Affinity-purified IgG (APIgG), clear liquid.

Host Animal:

Sheep

Immunogen:

Thrombin prepared from purified rabbit prothrombin, active site blocked with PPACK.

Concentration:

APIgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E^{1\%}_{280}$) of 13.4.

Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between – 10 and –20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for rabbit thrombin (or prothrombin) as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to rabbit thrombin (and prothrombin).

Neutralizing activity:

Not determined

Species Cross Reactivity: (immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX

Related Products:

Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantibility or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by: **AFFINITY BIOLOGICALS INC.** 1348 Sandhill Drive Ancaster ON CANADA L9G 4V5 Tel: (905) 304-9896 (800) 903-6020 Fax: (905) 304-9897 info@affinitybiologicals.com