

****REPRESENTATIVE DATASHEET******Sheep anti-human Tissue Factor (TF)**

Whole IgG from antiserum

10 mg

Product #: SATF-IG**Lot #:** XXXX**Expiry date:** XXXX

Store at -10 to -20°C

1348 Sandhill Drive, Ancaster, Ontario, Canada L9G 4V5
905-304-9896 • 800-903-6020 • fax 905-304-9897

For Research Use Only.

Not for use in diagnostic procedures.

Description of Tissue Factor (TF)

Tissue Factor (TF) is an integral membrane glycoprotein expressed in the plasma membranes of many cell types. It is a single chain molecule of 44 kDa consisting of an extra-cellular domain (residues 1-219), a trans-membrane domain (residues 220-242) and the C-terminal intracellular domain of residues 243-263. Most abundant in the tissue adventitia, TF becomes exposed to blood at the site of vascular injury. The availability of TF is important in initiating coagulation by acting as a receptor for both the zymogen and protease forms of plasma factor VII (FVII and FVIIa), as well as mediating the conversion of bound FVII to FVIIa. The binding of FVII to TF in the presence of a negatively charged surface such as a phospholipid (or cell surface) promotes the auto activation of F.VII by VIIa. The TF-FVIIa complex in the presence of calcium ions proteolytically activates factors IX and X. These enzyme products are then capable of activating FVII to FVIIa by feedback amplification. The activity of TF-FVIIa activity is regulated by a TFPI (tissue factor pathway inhibitor), a member of the Kunin superfamily of protease inhibitors. TFPI contains three kunitz domains and is able to bind and inhibit the TF-FVIIa complex in the presence of activated factor X and calcium ions. Antithrombin has also been reported to inhibit F.VIIa activity in the presence of TF and heparin. Although a membrane protein, low levels of TF products have been demonstrated in plasma. Increased levels of circulating TF products may be a risk factor for thrombotic disease¹⁻⁴.

REFERENCES and REVIEWS

1. Nemerson Y; in Hemostasis and Thrombosis, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 81-93, J.B. Lippincott Co., Philadelphia, 1994.
2. Neuenschwander PF, Morrissey JH; Deletion of the Membrane Anchoring Region of Tissue Factor Abolishes Autoactivation of F.VII but not Cofactor Function. JBC 267, pp 14477-14482, 1992.
3. Lawson LH, Butenas S, Mann KG; The Evaluation of Complex-dependent Alterations in Human Factor VIIa. JBC 267, pp 4834-4843, 1992.
4. Sambola A, Osende J, Hathcock J, Degen M, Nemerson Y, Fuster V, Crandall J, Badimon JJ; The Role of Risk Factors in the Modulation of Tissue Factor Activity and Blood Thrombogenicity. Circulation 107, pp 973-977, 2003.

Product Specifications**Description:**

Vial containing XXXX ml of whole IgG representing approximately 1 ml of antiserum. Total protein is 10 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Sheep

Immunogen:

Recombinant human tissue factor.

Concentration:IgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ($E^{1\%}_{280}$) of 13.4.**Buffer:**

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers.

Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for tissue factor as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of antibodies to human tissue factor.

Neutralizing activity:

Not determined.

Species Cross Reactivity:

Not determined.

Related Products:

Cat #: SATF-HRP	Sheep anti-human TF, IgG-peroxidase conjugate
Cat #: MATF-BIO	Murine anti-human TF, IgG-Biotin conjugate
Cat #: MATF-FITC	Murine anti-human TF, IgG-FITC conjugate

Visit our site (www.affinitybiologicals.com) for details.

Limited Warranty: This product is warranted to perform in accordance with its labeling and literature. Affinity Biologicals Inc. disclaims any implied warranty of merchantability or fitness for any other purposes, and in no event will Affinity Biologicals Inc. be liable for any consequential damages arising out of aforesaid express warranty.

Manufactured in Canada by:
AFFINITY BIOLOGICALS INC.
1348 Sandhill Drive
Ancaster ON
CANADA L9G 4V5
Tel: (905) 304-9896
(800) 903-6020
Fax: (905) 304-9897
info@affinitybiologicals.com

Document OPI0039, rev 3