


Datasheet



Mouse mAb to **MUC5AC/M1**
Clone **2-12M1**
Isotype **IgG1-κ**

Source

A BALB/c mouse was immunized with mucin isolated from an ovarian cyst fluid (pure endocervical type according to the Fenoglio's classification) from an ALe^b patient.
Fusion partner: Sp2/0.

Specifications

2-12M1 recognizes the peptide core of gastric mucin M1/MUC5AC, and more specifically with the 'c' epitope amongst the a, b, c, d, e, f, g, and h protein core epitopes defined by Bara for M1. 2-12M1 and 45M1 both specifically react with epitopes located in the C-terminal cysteine rich part of the peptide core of gastric mucin (MUC5AC). MUC5AC is present in primary ovarian mucinous cancer and gastric cancer, but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification pancreatic carcinoma and pre-cancerous changes vs. normal pancreas.




Figure 1: Human gastric mucosa stained for MUC5AC (paraffin)

Species reactivity

Positive: cat, cow, human, monkey, mouse.
Negative: rat.

Applications

2-12M1 reacts with MUC5AC after protease/periodate pretreatment. Immunoblotting: strongly positive without beta-mercaptoethanol pretreatment of mucin solution. ELISA or IRMA as component of a mixture of anti-M1 MAbs.

ELISA	Frozen sections	Paraffin sections	Western blot
+	+	Protease/periodate	+

Format

Produced in tissue culture, contains no host Ig. Antibodies are affinity purified and presented in PBS with 0,02 % sodium azide.

Stored at 4°C- 8°C, shelf life is at least 24 months after purchase.

Dilution advice

- ELISA (solid phase: 0,1-100 µg/ml; tracer: 0,001-100 µg/ml for 30 min at RT).
- Immunoblotting (1-2 µg/ml).
- Immunohistology (1-2 µg/ml for 30 min at RT; staining of formalin-fixed tissues requires 0,1% pronase digestion (0,1% CaCl₂, 0,1N NaOH) for 15min. at 37°C; staining is enhanced by subsequent incubation in 20mM periodic acid in 50mM acetate buffer pH 5.0 for 30 minutes at RT in the dark).

Positive control

Stomach, gastric cancer.

Datasheet



References

- Bara, J. et al., *Cancer Res.* **46**: 3983-3989 (1986).
- Bara, J. et al., *Biochem. J.* **254**: 185-193 (1988).
- Bara, J. et al., *Int. J. Cancer* **47**: 304-310 (1991).
- Bara, J. et al., *J. Immunol. Methods* **149**: 105-113 (1992).
- Guyonnet Duperat V. et al., *Biochem. J.* **305**: 211 219 (1995).