

Anti Chromogranin A (1-28)(Rat) Serum

Cat. No. Y294

Lot No. 0791101012

Description: This antiserum was raised in a rabbit by immunization with a conjugate of bovine serum albumin (BSA) to synthetic chromogranin A (1-28) (rat) peptide at position 17 Cys residue. The product vial contains 50 μ L of the titled antiserum obtained by lyophilizing its 0.001M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with chromogranin A (rat).

Immunogen: Synthetic Chromogranin A (1-28) (rat), BSA conjugate**Host:** Rabbit

1

28

Amino Acid Sequence of Chromogranin A (1-28)(rat)^{1,2)}: LPVNSPMTKG DTKVMKCVLE VISDSLK**Product Form:** Lyophilized unpurified serum**Size:** 50 μ L

Reconstitution: Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN₃ 0.1%).


Storage: The product will be stable for over one year if stored at -20 °C to -80 °C until opened. Upon reconstitution, the antiserum solution must be stored at 2 °C to 8 °C and used within one month. Repeated freezing- thawing should be avoided.

Suggested Working Dilution Range: 1:3,000 (final dilution 1:21,000) for radioimmunoassay^{3,4)}; 1:500~3,000 for immunohistochemistry (frozen section)³⁾. Optimal dilution should be determined by each laboratory for each application.

Specificity (based on RIA): Chromogranin A (1-28) (rat) 100%, Pancreastatin (rat) 0%.**Positive Control (immunohistochemistry):** Rat adrenal gland³⁾**Species Tested:** Rat**REFERENCES:**

- 1) A. Iacangelo, H. Okayama et al., primary structure of rat chromogranin A and distribution of its mRNA. FEBS Letter 227: 115-121, 1988
- 2) R.J. Parmer. A.H. Koop et al., Molecular cloning of chromogranin A from rat pheochromocytoma cells. Hypertension 14: 435-444, 1989
- 3) K. Ohshima, S. Nagasawa et al., Synthesis of rat chromogranin A fragments and their immunological properties. Peptide Chemistry, 1991, Protein Research Foundation, Osaka, A. Suzuki (Ed), pp157-162, 1992
- 4) S. Nagasawa, Y. Nishikawa et al., Distribution of chromogranin A-like peptides in the rat. Biomedical Research 16 (2): 83-90, 1995

FOR RESEARCH LABORATORY USE ONLY
DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM



Yanaihara Institute Inc.

2480-1 Awakura, Fujinomiya, Shizuoka, 418-0011 Japan
TEL: +81-544-22-2771 FAX: +81-544-22-2770

For more information, please visit our internet website: <http://www.yanaihara.co.jp> or contact us to: ask@yanaihara.co.jp