

Rat Anti-Murine Factor XIII

Clone GMA-741

Plasma factor XIII is a tetrameric molecule composed of two A subunits (83kDa) and two B subunits (80kDa). Thrombin cleaves a peptide bond within the A chain to form activated factor XIII. Factor XIIIa is the final component of the blood clotting cascade and is responsible for crosslinking fibrin. GMA-741 binds murine Factor XIII in solid-phase ELISA. It does not cross-react with human Factor XIII.

Description		
Antibody Source:	rat monoclonal, IgG2 _{aK}	
Antigen Species Bound:	mouse	
Specificity:	murine Factor XIII	
Immunogen:	murine Factor XIII	

Formulation and Storage	
Purity:	Purified by protein G affinity chromatography from serum-free cell culture supernatant.
Product Formulation:	Lyophilized from a ≥ 1 mg/ml solution in 20 mM NaH $_2$ PO $_4$ 0.15 M NaCl, 1.0% (w/v) mannitol, pH 7.4. Concentration determined by absorbance measurement at 280 nm and using an extinction coefficient of 1.4 ($\epsilon_{0.1\%}$).
Reconstitution:	Reconstitute with deionized water.
Storage:	Store lyophilized or reconstituted and aliquoted material at -20 °C for prolonged periods. Avoid freeze-thaw cycles. Alternatively, add 0.02% (w/v) sodium azide to reconstituted solution and store at 4 °C.
Country of origin:	USA
Size Options:	0.1 mg or 0.5 mg

Applications	
Working Concentration:	Approximately 1-5 µg/ml. Researcher should titer antibody in specific assay.
ELISA:	Binds murine Factor XIII in solid-phase ELISA.
Immunoblotting:	Not recommended.

